

The writer uses a table to summarize the findings presented in two sources.

Table 1
Effectiveness of Sibutramine and Orlistat in Adolescents

Medication	Subjects	Treatment ^a	Side effects	Average weight loss/gain
Sibutramine	Control	0-6 mos.: placebo 6-12 mos.: sibutramine	Mos. 6-12: increased blood pressure; increased pulse rate	After 6 mos.: loss of 3.2 kg (7 lb) After 12 mos.: loss of 4.5 kg (9.9 lb)
	Medicated	0-12 mos.: sibutramine	Increased blood pressure; increased pulse rate	After 6 mos.: loss of 7.8 kg (17.2 lb) After 12 mos.: loss of 7.0 kg (15.4 lb)
Orlistat	Control	0-12 mos.: placebo	None	Gain of 0.67 kg (1.5 lb)
	Medicated	0-12 mos.: orlistat	Oily spotting; flatulence; abdominal discomfort	Loss of 1.3 kg (2.9 lb)

A note gives the source of the data.

Note. The data on sibutramine are adapted from "Behavior Therapy and Sibutramine for the Treatment of Adolescent Obesity," by R. I. Berkowitz, T. A. Wadden, A. M. Tershakovec, & J. L. Cronquist, 2003, *Journal of the American Medical Association*, 289, pp. 1807-1809. The data on orlistat are adapted from *Xenical (Orlistat) Capsules: Complete Product Information*, by Roche Laboratories, December 2003, retrieved from <http://www.rocheusa.com/products/xenical/pi.pdf>

A content note explains data common to all subjects.

^aThe medication and/or placebo were combined with behavioral therapy in all groups over all time periods.